

AD-A001 936

MASSACHUSETTS INST OF TECH CAMBRIDGE DEPT OF CHEMISTRY F/6 7/3
NEW FUNCTIONAL ALLYLIC LITHIUM REAGENTS: 6EM-DIALKOXYALLYLLITHI--ETC(U)
1979 D SEYFERTH, R E MAMMARELLA, H A KLEIN N00014-76-C-0037

UNCLASSIFIED

NL

1-1
7-5-77

■



END
DATE
FILMED
4-80
DTIC



LEVEL II

(12)

NEW FUNCTIONAL ALLYLIC LITHIUM REAGENTS: GEM-DIALKOXYALLYLLITHIUM REAGENTS: A USEFUL ROUTE TO β -SILYL AND β -STANNYLPROPIONATE ESTERS

(10) Dietmar Seyferth / Robert E. Mammarella and Helmut A. Klein

Department of Chemistry
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

DTIC
ELECT
MAR 14 1980
S C

1, 1749 /
(12) 121
ABSTRACT

The reaction of sec-butyllithium with acrolein dialkyl acetals in THF or in THF/Et₂O/pentane at -95°C results in formation of gem-dialkoxyallyllithium reagents, $\text{Li}[\text{CH}_2\text{CHC}(\text{OR})_2]$. These react with organosilicon and organotin chlorides to give ketene acetals, $\text{R}_3\text{SiCH}_2\text{CH}=\text{C}(\text{OR})_2$ and $\text{R}_3\text{SnCH}_2\text{CH}=\text{C}(\text{OR})_2$. The acid hydrolysis of these products produces β -substituted propionic acid esters, $\text{R}_3\text{SiCH}_2\text{CH}_2\text{CO}_2\text{R}$ and $\text{R}_3\text{SnCH}_2\text{CH}_2\text{CO}_2\text{R}$. Reactions of these lithium reagents with allyl bromide gave esters of 5-hexenoic acid, $\text{CH}_2=\text{CH}(\text{CH}_2)_3\text{CO}_2\text{R}$ (R = Me, Et).

(15)
Contract / N00014-76-C-0837 /

This document has been approved for public release and sale; its distribution is unlimited.

ADA 081936




DOC FILE COPY

220 007

30 2 19 076

INTRODUCTION


The lithiation of allylic ethers provides useful reagents which are operationally equivalent to a β -acyl carbanion (eq. 1-3)¹.



In this context, the possible lithiation of an allylic acetal, $\text{CH}_2=\text{CHCH(OR)}_2$, was of some interest. The expected product is a gem-dialkoxyallyllithium reagent (eq. 4) which potentially could react



with an electrophile, E, at either the unsubstituted or the substituted terminus of the allyl group (eq. 5). If the new bond is formed



at the substituted end, an allylic acetal would be formed. Hydrolysis of the latter would give a vinyl ketone. In this case the reagent would function as a carbonyl anion equivalent, $\text{CH}_2=\text{CHC(O)}^-$. If, on the other hand, the new bond is formed at the unsubstituted terminus of the allyl group, a γ -substituted ketene acetal would result. Hydrolysis of such a product would give an ester, as shown in eq. 5. In

this case the reagent would be operationally equivalent to a β -alkoxycarbonyl carbanion, $\text{ROC(O)CH}_2\text{CH}_2^-$.

Of special interest to us were the reactions of such $\text{Li} \cdot [\text{CH}_2\text{CHC(OR)}_2]$ reagents with organometallic halides, in particular trimethylchlorosilane and trimethyltin chloride. Alkyl- and aryl-substituted allylic lithium reagents of type $\text{Li} [\text{CH}_2\text{CHCHR}]$ and $\text{Li} [\text{CH}_2\text{CHCR}_2]$ had been found to react with both of these halides to give exclusively products of structure $\text{Me}_3\text{MCH}_2\text{CH=CHR}$ and $\text{Me}_3\text{MCH}_2\text{CH=CR}_2$ ($\text{M} = \text{Si}$ and Sn), respectively.^{2,3} On the other hand, trimethylchlorosilane reacted with gem-dichloro-⁴ and gem-difluoroallyllithium⁵ to give $\text{Me}_3\text{SiCX}_2\text{CH=CH}_2$ ($\text{X} = \text{Cl}$ and F). Trimethyltin chloride formed $\text{Me}_3\text{SnCH}_2\text{CH=CCl}_2$ on reaction with $\text{Li} [\text{CH}_2\text{CHCCl}_2]$. In the case of the Me_3SiCl reactions it was quite likely that products of kinetic control were obtained, while those derived from Me_3SnCl were products of thermodynamic control.⁴


In view of the potential synthetic utility of reagents of $\text{Li} [\text{CH}_2\text{CHC(OR)}_2]$ and in view of the uncertainty of the regioselectivity in their reactions with various electrophiles, we have investigated their synthesis, stability and, to a limited extent, their reactions.

RESULTS AND DISCUSSION

We have found that acrolein dimethyl and diethyl acetals may be lithiated by sec-butyllithium in a THF/diethyl ether/pentane solvent mixture at $-90^\circ \pm 5^\circ\text{C}$ to give a yellow solution containing the respective gem-dialkoxyallyllithium reagents (eq. 4, $\text{R} = \text{Me}$ and Et ; $\text{R}' = \text{sec-C}_4\text{H}_9$). The temperature range for successful preparation of these reagents is quite narrow: below -100°C the metalation reaction appears to be too slow; above -85°C the reagents begin to decompose slowly. Their decomposition is rapid at -65°C and above. Nonethe-


less, they find useful applications in synthesis.

We have thus far studied the reactions of these new allylic lithium reagents with several organosilicon and organotin chlorides and with allyl bromide (Table 1). Such reactions of $\text{Li}[\text{CH}_2\text{CHC(OR)}_2]$ with organosilicon and organotin chlorides, followed by nonhydrolytic work-up, gave silicon- and tin-substituted ketene acetals as the exclusive products (eq. 6). These acetals are acid-sensitive and



decompose slowly on standing.

The reaction shown in eq. 6 thus provides a useful new route to substituted ketene-O,O-acetals. Such reactions of gem-dialkoxyallyllithium reagents with silicon and tin chlorides, when followed by hydrolysis of the initially formed products, ^{provide} a new synthesis of β -silyl- and β -stannylpropionic acid esters as shown in eq 7. The



yields of these products ranged from moderate to excellent. The conditions required for hydrolysis of the initially produced acetals differed for the reactions with organosilicon and organotin chlorides. In the case of the former, hydrolysis with neutral water was sufficient to give complete conversion to the esters. The organotin-substituted acetals were only incompletely hydrolyzed on treatment of the reaction mixtures with neutral water. Hydrolysis with 5%, or better, 10%, aqueous hydrochloric acid was required for production of the tin-substituted ester. The explanation for these observations is a simple one. Hydrolysis of unreacted chlorosilane, generally added in excess, generates an acid medium which facilitates hydrolysis of the

TABLE 1. Reactions of gem-Dialkoxyallyllithium Reagents^a with Chlorosilanes, Tin Chlorides and Allyl Bromide.

Acrolein Dialkyl Acetal (mmol)	Substrate (mmol)	Work-up	Product	(% Yield) ^d
$\text{CH}_2=\text{CHCH}(\text{OMe})_2$ (21)	Me_3SiCl (27.6)	5% HCl	$\text{Me}_3\text{SiCH}_2\text{CH}_2\text{CO}_2\text{Me}$	(75)
$\text{CH}_2=\text{CHCH}(\text{OMe})_2$ (42)	Me_3SiCl (45	anhydrous	$\text{Me}_3\text{SiCH}_2\text{CH}=\text{C}(\text{OMe})_2$	(70) ^b
$\text{CH}_2=\text{CHCH}(\text{OMe})_2$ (43)	Me_2SiCl_2 (21)	5% HCl	$\text{Me}_2\text{Si}(\text{CH}_2\text{CH}_2\text{CO}_2\text{Me})_2$	(61)
$\text{CH}_2=\text{CHCH}(\text{OMe})_2$ (21)	Et_3SiCl (21)	5% HCl	$\text{Et}_3\text{SiCH}_2\text{CH}_2\text{CO}_2\text{Me}$	(70) ^c
$\text{CH}_2=\text{CHCH}(\text{OMe})_2$ (42)	PhMe_2SiCl (49)	5% HCl	$\text{PhMe}_2\text{SiCH}_2\text{CH}_2\text{CO}_2\text{Me}$	(77)
$\text{CH}_2=\text{CHCH}(\text{OEt})_2$ (25)	Me_3SiCl (40)	H_2O	$\text{Me}_3\text{SiCH}_2\text{CH}_2\text{CO}_2\text{Et}$	(43)
$\text{CH}_2=\text{CHCH}(\text{OEt})_2$ (63)	Et_3SiCl (86)	H_2O	$\text{Et}_3\text{SiCH}_2\text{CH}_2\text{CO}_2\text{Et}$	(86)(71% distilled)
$\text{CH}_2=\text{CHCH}(\text{OEt})_2$ (63)	PhMe_2SiCl (86)	H_2O	$\text{PhMe}_2\text{SiCH}_2\text{CH}_2\text{CO}_2\text{Et}$	(53)
$\text{CH}_2=\text{CHCH}(\text{OMe})_2$ (42)	Me_3SnCl (60)	10% HCl	$\text{Me}_3\text{SnCH}_2\text{CH}_2\text{CO}_2\text{Me}$	(72)
$\text{CH}_2=\text{CHCH}(\text{OMe})_2$ (42)	Me_3SnCl (60)	anhydrous	$\text{Me}_3\text{SnCH}_2\text{CH}=\text{C}(\text{OMe})_2$	(69)
$\text{CH}_2=\text{CHCH}(\text{OEt})_2$ (25)	Me_3SnCl (40)	5% HCl	$\text{Me}_3\text{SnCH}_2\text{CH}_2\text{CO}_2\text{Et}$	(27)
$\text{CH}_2=\text{CHCH}(\text{OEt})_2$ (25)	Me_3SnCl (40)	H_2O	$\text{Me}_3\text{SnCH}_2\text{CH}_2\text{CO}_2\text{Et}$ (1 part) + $\text{Me}_3\text{SnCH}_2\text{CH}=\text{C}(\text{OEt})_2$ (2 parts)	
$\text{CH}_2=\text{CHCH}(\text{OEt})_2$ (25)	Et_3SnCl (32)	5% HCl	$\text{Et}_3\text{SnCH}_2\text{CH}_2\text{CO}_2\text{Et}$	(24)
$\text{CH}_2=\text{CHCH}(\text{OMe})_2$ (42)	$\text{CH}_2=\text{CHCH}_2\text{Br}$ (60)	5% HCl	$\text{CH}_2=\text{CHCH}_2\text{CH}_2\text{CH}_2\text{CO}_2\text{Me}$	(47)
$\text{CH}_2=\text{CHCH}(\text{OEt})_2$ (25)	$\text{CH}_2=\text{CHCH}_2\text{Br}$ (40)	5% HCl	$\text{CH}_2=\text{CHCH}_2\text{CH}_2\text{CH}_2\text{CO}_2\text{Et}$	(22)

TABLE 1: Footnotes

- ^a Prepared at -90° 5°C by reaction with sec-butyllithium.
- ^b A 12:1 mixture of $\text{Me}_3\text{SiCH}_2\text{CH}=\text{C}(\text{OMe})_2$ and $\text{Me}_3\text{SiCH}_2\text{CH}_2\text{CO}_2\text{Me}$ was produced.
- ^c The $\text{Et}_3\text{SiOSiEt}_3$ formed on hydrolysis of unconverted Et_3SiCl cannot be separated from the product and so it is better to use an excess of the lithium reagent.
- ^d By-products included the respective disiloxane in those reactions in which an excess of the chlorosilane was used.

acetal. Organotin chlorides, on the other hand, do not hydrolyze readily. Although their rate of hydrolysis is rapid, the hydrolysis equilibrium is not favorable.⁶ Hence, insufficient acid is generated and external acid must be added to effect hydrolysis of the acetal.

The formation of products of type $R'_3MCH_2CH=C(OR)_2$ rather than $R'_3MC(OR)_2CH=CH_2$ in these reactions of $Li[CH_2CHC(OR)_2]$ with R'_3MCl may be understood in terms of the operation of steric effects. It is unlikely that allylic organosilicon compounds of either type ($M = Si$ in the formulas above) would take part in transmetalation equilibria, so we assume that we are dealing with products of kinetic control. Products of type $R'_3SiC(OR)_2CH=CH_2$ would have been of some interest: their hydrolysis should have given α -silyl ketones, $R'_3SiC(O)CH=CH_2$.

Other methods for the preparation of β -silyl- and β -stannyl-propionic acid esters are available. Such silyl-substituted esters can be prepared by the acetoacetic ester and malonic ester syntheses when the appropriate halomethylsilanes, R_3SiCH_2X , are available.⁹ Both silyl and stannyl-substituted esters have been prepared by silicon or tin hydride addition to acrylate esters,¹⁰⁻¹³ while the reaction ^{of} β -halopropionic acid esters with tin foil provided another route to β -stannyl-substituted esters.¹⁴ Thus the present procedure, which is based on the availability of the silicon or tin halide, complements these procedures.

EXPERIMENTAL

General Comments.

All reactions were carried out in flame-dried, nitrogen-flushed glassware under an atmosphere of prepurified nitrogen or argon. Solvents were rigorously dried prior to use. The reaction temperatures which are reported are uncorrected. They were obtained using a pentane total immersion thermometer which was immersed to a depth of about 3 cm into the stirred reaction mixture. Experiments showed that a stem correction of -8° to -10°C is appropriate.

sec-Butyllithium was purchased from Alfa Products, Thiokol/Ventron Corp., chlorosilanes from Petrarch Systems, Inc., the acrolein dialkyl acetals from Aldrich Chemical. Co. Methyltin starting materials were kindly donated by Cincinnati Milacron Chemicals, Inc.

Infrared spectra were recorded using a Perkin Elmer Model 457A grating infrared spectrophotometer, NMR spectra with a Varian Associates T60 spectrometer. Gas-liquid chromatography (GLC) was employed for analysis of reaction products, isolation of product samples and for yield determination by the internal standard method.

Preparation of gem-Dialkoxallyllithium Reagents: General Procedure.

A Morton (creased) flask of appropriate size, equipped with a paddle-type stirrer, a no-air stopper and a Claisen adapter which was fitted with a low temperature thermometer and a gas inlet tube, was flame-dried and then allowed to cool under a stream of nitrogen or argon. The flask then was charged with the solvent mixture (75 ml of THF, 15 ml of diethyl ether and 15 ml of pentane), 21 mmol of the acrolein dialkyl acetal (2.5 ml in the case of $\text{CH}_2=\text{CHCH}(\text{OMe})_2$) was added, and the solution was cooled to $-90^\circ \pm 5^\circ\text{C}$ by partial immersion in a liquid nitrogen Dewar flask. To this solution was added with stirring and under nitrogen (or argon) 20 ml of 1.1M sec-butyl-lithium in cyclohexane (22 mmol). The temperature was maintained at $-90^\circ \pm 5^\circ\text{C}$ during the course of this very slow, dropwise addition. The resulting yellow solution was stirred 2-3 hr. at -90°C .

To this reagent solution then was added the chlorosilane or the tin chloride (slight excess, ca. 25-30 mmol) by syringe over a 10 min. period. The resulting mixture, now milky white, was stirred at -90°C for 30 min. and then was allowed to warm to room temperature.

In the non-hydrolytic work-up, the reaction mixture was transferred by cannula to a dry distillation flask. Solvents were removed at reduced pressure and the residue was trap-to-trap distilled at ca. 0.1 mm Hg into a flask cooled with liquid nitrogen. The distillate then was examined by GLC.

In the hydrolytic work-up, the cannulated solution was poured into a separatory funnel and treated successively with distilled water, two portions of 5 or 10% hydrochloric acid and, again, water. The aqueous phases were back-extracted with pentane and the combined organic phases were dried (MgSO_4) and the solvents removed at

reduced pressure. The residue was trap-to-trap distilled, as above.

Some reactions were carried out on a larger scale (42 mmol of $\text{CH}_2=\text{CHCH}(\text{OMe})_2$, 63 mmol of $\text{CH}_2=\text{CHCH}(\text{OEt})_2$). In some cases THF alone, rather than the mixed solvent system, was used, but the mixed solvent is preferred.

Organosilicon and Organotin Products: Characterization.

Table 2 lists the organosilicon and organotin esters and acetals prepared, their refractive indices, analyses and spectroscopic properties. The infrared spectra of the esters (liquid film) showed strong ester carbonyl absorption at $1735\text{--}1740\text{ cm}^{-1}$. The ketene acetals showed strong bands in their IR spectra at 1675 cm^{-1} ($\text{Me}_3\text{SiCH}_2\text{CH}=\text{C}(\text{OMe})_2$) and 1670 cm^{-1} ($\text{Me}_3\text{SnCH}_2\text{CH}=\text{C}(\text{OMe})_2$).

The methyl and ethyl esters of 5-hexenoic acid are known compounds.

$\text{CH}_2=\text{CHCH}_2\text{CH}_2\text{CH}_2\text{CO}_2\text{Me}$, n_D^{20} 1.4205, a known compound.⁷

IR (film): $\nu(\text{C}=\text{O})$ 1755, $\nu(\text{C}=\text{C})$ 1640 cm^{-1} .

NMR (CDCl_3): δ 1.6-2.5 (m, 6H, $\text{CH}_2\text{CH}_2\text{CH}_2$), 3.6 (s, 3H, CH_3) and 4.75-6.3 (m, 3H, $\text{CH}=\text{CH}_2$).

$\text{CH}_2=\text{CHCH}_2\text{CH}_2\text{CH}_2\text{CO}_2\text{Et}$, n_D^{20} 1.4225; lit.⁸ n_D^{20} 1.4220

IR (film): $\nu(\text{C}=\text{O})$ 1735, $\nu(\text{C}=\text{C})$ 1640 cm^{-1} .

NMR (CCl_4): δ 1.22 (t, 3H, J 7Hz, CH_3), 1.48-2.03 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2$), 2.10 and 2.22 (2t, 4H, J 7Hz, $\text{CH}_2\text{CH}_2\text{CH}_2$); 4.03 (q, 2H, OCH_2), 4.73-6.17 (m, 3H, $\text{CH}=\text{CH}_2$).

Acknowledgments. This work was supported in part by the Office of Naval Research. H.A.K. is grateful to the University of Kiel for a leave of absence and to the Max Kade Foundation for the award of a postdoctoral fellowship.

TABLE 1. Reaction Products.

Compound	n_D^{20} ($^{\circ}\text{C}$)	Analysis, %		^1H NMR δ (ppm)
		Found	(Calcd.)	
		Carbon	Hydrogen	
$\text{Me}_3\text{SiCH}_2\text{CH}_2\text{CO}_2\text{CH}_3$	1.4190 (20) ^a	52.42 (52.45)	10.04 (10.06)	0.00 (s, 9H, Me_3Si), 0.63-1.00 (m, 2H, SiCH_2), 2.1-2.44 (m, 2H, CH_2CO), 3.6 (s, 3H, OCH_3) (in CDCl_3)
$\text{Me}_2\text{Si}(\text{CH}_2\text{CH}_2\text{CO}_2\text{Me})_2$	1.4396 (25)	51.25 (51.22)	8.60 (8.62)	0.00 (s, 6H, Me_2Si), 0.65-1.05 (m, 4H, SiCH_2), 2.1-2.5 (m, 4H, CH_2CO), 3.6 (s, 6H, OCH_3) (in CDCl_3)
$\text{Et}_3\text{SiCH}_2\text{CH}_2\text{CO}_2\text{CH}_3$	1.4409 (25)	59.19 (59.40)	10.67 (10.89)	0.15-1.25 (m, 17H, Et_3Si and SiCH_2), 1.95-2.4 (m, 2H, CH_2CO), 3.55 (s, 3H, OCH_3) (in CDCl_3)
$\text{PhMe}_2\text{SiCH}_2\text{CH}_2\text{CO}_2\text{CH}_3$	1.4950 (25)	64.74 (64.86)	8.21 (8.11)	0.2 (s, 6H, SiMe_2), 0.8-1.2 (m, 2H, SiCH_2), 2.0-2.4 (m, 2H, CH_2CO), 3.45 (s, 3H, OCH_3), 7.2-7.5 (m, 5H, Ph) (in CDCl_3)
$\text{Me}_3\text{SiCH}_2\text{CH}_2\text{CO}_2\text{C}_2\text{H}_5$	b	54.91 (55.12)	10.40 (10.41)	0.00 (s, 9H, Me_3Si), 0.62-1.02 (m, 2H, SiCH_2), 1.23 (t, J 7Hz, 3H, CH_3 of Et), 2.02-2.37 (m, 2H, CH_2CO), 4.02 (q, J 7Hz, CH_2 of Et) (in CCl_4)
$\text{Et}_3\text{SiCH}_2\text{CH}_2\text{CO}_2\text{C}_2\text{H}_5$	1.4407 (25) ^c	61.04 (61.05)	11.12 (11.18)	0.27-1.18 (m, 17H, Et_3Si and SiCH_2), 1.27 (t, J 7Hz, 3H, CH_3 of OEt), 2.03-2.38 (m, 2H, CH_2CO), 4.05 (q,

J 7Hz, 2H, OCH₂)

PhMe ₂ SiCH ₂ CH ₂ CO ₂ C ₂ H ₅	1.4917 (25) ^d	66.09 (66.05)	8.52 (8.53)	0.32 (s, 6H, Me ₂ Si), 1.05 (t, J 7Hz, 2H, SiCH ₂), 1.33 (t, J 7Hz, 3H, CH ₃ of OEt), 2.08-2.42 (m, 2H, CH ₂ CO), 4.03 (q, 2H, CH ₂ of OEt), 6.88-7.57 (m, 5H, Ph) (in CCl ₄)
Me ₃ SnCH ₂ CH ₂ CO ₂ CH ₃	1.4695 (20) ^e	33.44 (33.50)	6.49 (6.38)	0.07 (s, 9H, Me ₃ Sn, J(¹¹⁹ Sn- ¹ H) 52 Hz, J(¹¹⁷ H- ¹ H) 50 Hz), 1.00 (t, J 8Hz, 2H, SnCH ₂), 2.50 (t, J 8Hz, 2H, CH ₂ CO), 3.63 (s, 3H, OCH ₃) (in CDCl ₃)
Me ₃ SnCH ₂ CH ₂ CO ₂ C ₂ H ₅	1.4638 (25) ^f	36.64 (36.27)	6.92 (6.85)	0.08 (s, 9H, Me ₃ Sn, J(¹¹⁹ Sn- ¹ H) 52 Hz, J(¹¹⁷ H- ¹ H) 50 Hz), 0.95 (t, 7Hz, 2H, SnCH ₂), 1.27 (t, J 7Hz, 3H, CH ₃ of OEt), 2.50 (t, J 7Hz, 2H, CH ₂ CO), 4.08 (q, 2H, OCH ₂) (in CCl ₄)
Et ₃ SnCH ₂ CH ₂ CO ₂ C ₂ H ₅	1.4760 (25) ^g	43.51 (43.04)	7.90 (7.88)	0.33-1.48 (m, 20H, Et ₃ Sn and CH ₃ of OEt), 2.50 (t, 7Hz, CH ₂ CO), 4.08 (q, J 7Hz, 2H, OCH ₂) (in CCl ₄)
Me ₃ SiCH ₂ CH=C(OMe) ₂	1.4354 (20)	55.15 (55.17)	10.70 (10.34)	0.00 (s, 9H, Me ₃ Si), 1.28 (d, J 8Hz, 2H, SiCH ₂), 3.13-3.81 (m, 1H, =CH), 3.50 and 3.56 (two s, each 3H, OCH ₃) (in CDCl ₃)

$\text{Me}_3\text{SnCH}_2\text{CH}=\text{C}(\text{OMe})_2$ 1.4845 (20) 36.15 6.82 0.05 (s, 9H, Me_3Sn , $J(^{119}\text{Sn}-^1\text{H})$ 52 Hz, $J(^{117}\text{Sn}-^1\text{H})$ 50 Hz), 1.52 (d, J 8 Hz, 2H, SnCH_2), 3.15-3.90 (m, 1H, =CH), 3.46 and 3.53 (two s, each 3H, OCH_3) (in CDCl_3)

$\text{Me}_3\text{SnCH}_2\text{CH}=\text{C}(\text{OEt})_2$ 1.4640 (25) 40.90 7.65 0.07 (s, 9H, Me_3Sn , $J(^{119}\text{Sn}-^1\text{H})$ 52 Hz, $J(^{117}\text{Sn}-^1\text{H})$ 50 Hz), 1.25 and 1.28 (two t, J 7 Hz, each 3H, CH_3 of OEt), 1.52 (d, J 8 Hz, 2H, SnCH_2), 3.42 (m, 5H, OCH_2 and =CH) (in CCl_4)

a Lit. 15 $n^{20}\text{D}$ 1.4180; b Lit. 9 $n^{20}\text{D}$ 1.4198; lit. 16 $n^{20}\text{D}$ 1.4205; c Lit. 12 $n^{20}\text{D}$ 1.4422; d Lit. 9 $n^{20}\text{D}$ 1.4972; e Lit. 14 $n^{20}\text{D}$ 1.4690; f Lit. 14 $n^{20}\text{D}$ 1.4660; g Lit. 12 $n^{20}\text{D}$ 1.4762.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	<i>for info</i>
By	
Distribution/	
Availability Codes	
Dist.	Avail and/or special
<i>A</i>	

REFERENCES

1. (a) D. A. Evans, G. C. Andrews and R. Buckwalter, J. Amer. Chem. Soc., 96 (1974) 5560; (b) W. C. Still and T. L. Macdonald, J. Amer. Chem. Soc., 96 (1974) 5561; (c) W. C. Still and T. L. Macdonald, J. Org. Chem., 41 (1976) 3620; (d) A. P. Kozikowski and K. Isobe, Tetrahedron Lett. (1979) 833.
2. D. Seyferth and T. F. Jula, J. Organometal. Chem., 66 (1974) 195
3. D. Seyferth and R. E. Mammarella, J. Organometal. Chem., 177 (1979) 53.
4. D. Seyferth, G. J. Murphy and R. A. Woodruff, J. Organometal. Chem., 141 (1977) 71.
5. D. Seyferth and K. R. Wursthorn, J. Organometal. Chem., 137 (1977) C17.
6. R. H. Prince, J. Chem. Soc. (1959) 1783.
7. P. Baudart, Bull. Soc. Chim. France (1946) 86.
8. Yu. N. Ogibin, M. I. Katzin and G. I. Nikishin, Synthesis (1974) 889.
9. L. H. Sommer and N. S. Marans, J. Amer. Chem. Soc., 72 (1950) 1935.
10. J. L. Speier, J. A. Webster and G. H. Barnes, J. Amer. Chem. Soc., 79 (1957) 974.
11. G. J. M. Van der Kerk and J. G. Noltes, J. Appl. Chem., 9 (1959) 106.
12. F. Rijks, M. J. Janssen, W. Drenth and G. J. M. Van der Kerk, J. Organometal. Chem., 2 (1964) 347.
13. R. E. Hutton, J. W. Burley and V. Oakes, J. Organometal. Chem., 156 (1978) 369.
14. M. Nomuri, N. Matsui and S. Matsuda, Kogyo Kagaku Zasshi, 71 (1968) 1526; Chem. Abstr., 70 (1969) 47572.

15. A. D. Petrov, S. I. Sadykh-Zade and E. I. Filatova, Zh. Obshch. Khim., 29 (1959) 2936.
16. N. V. Komarov and N. V. Semenova, Zh. Obshch. Khim., 36 (1966) 103.

TECHNICAL REPORT DISTRIBUTION LIST, 053

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Dr. R. N. Grimes University of Virginia Department of Chemistry Charlottesville, Virginia 22901	1	Dr. M. H. Chisholm Department of Chemistry Indiana University Bloomington, Indiana 47401	1
Dr. M. Tsutsui Texas A&M University Department of Chemistry College Station, Texas 77843	1	Dr. B. Foxman Brandeis University Department of Chemistry Waltham, Massachusetts 02154	1
Dr. M. F. Hawthorne University of California Department of Chemistry Los Angeles, California 90024	1	Dr. T. Marks Northwestern University Department of Chemistry Evanston, Illinois 60201	1
Dr. D. B. Brown University of Vermont Department of Chemistry Burlington, Vermont 05401	1	Dr. G. Geoffrey Pennsylvania State University Department of Chemistry University Park, Pennsylvania 16802	1
Dr. W. B. Fox Naval Research Laboratory Chemistry Division Code 6130 Washington, D.C. 20375	1	Dr. J. Zuckerman University of Oklahoma Department of Chemistry Norman, Oklahoma 73019	1
Dr. J. Adcock University of Tennessee Department of Chemistry Knoxville, Tennessee 37916	1	Professor O. T. Beachley Department of Chemistry State University of New York Buffalo, New York 14214	1
Dr. A. Cowley University of Texas Department of Chemistry Austin, Texas 78712	1	Professor P. S. Skell Department of Chemistry The Pennsylvania State University University Park, Pennsylvania 16802	1
Dr. W. Hatfield University of North Carolina Department of Chemistry Chapel Hill, North Carolina 27514	1	Professor K. M. Nicholas Department of Chemistry Boston College Chestnut Hill, Massachusetts 02167	1
Professor H. Abrahamson University of Oklahoma Department of Chemistry Norman, Oklahoma 73019	1	Professor R. Neilson Department of Chemistry Texas Christian University Fort Worth, Texas 76129	1
		Professor M. Newcomb Texas A&M University Department of Chemistry College Station, Texas 77843	1

TECHNICAL REPORT DISTRIBUTION LIST, GENNo.
Copies

Dr. Rudolph J. Marcus
Office of Naval Research
Scientific Liaison Group
American Embassy
APO San Francisco 96503

1

Mr. James Kelley
DTNSRDC Code 2803
Annapolis, Maryland 21402

1

Nov. 1979

list

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Office of Naval Research Attn: Code 472 800 North Quincy Street Arlington, Virginia 22217	2	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 1211 Research Triangle Park, N.C. 27709	1
ONR Branch Office Attn: Dr. George Sandoz 536 S. Clark Street Chicago, Illinois 60605	1	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
ONR Branch Office Attn: Scientific Dept. 715 Broadway New York, New York 10003	1	Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555	1
ONR Branch Office 1030 East Green Street Pasadena, California 91106	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	-
ONR Branch Office Attn: Dr. L. H. Peebles Building 114, Section D 666 Summer Street Boston, Massachusetts 02210	1	Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Director, Naval Research Laboratory Attn: Code 6100 Washington, D.C. 20390	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
The Assistant Secretary of the Navy (R,E&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Office of Naval Research Attn: Dr. Richard S. Miller 800 N. Quincy Street Arlington, Virginia 22217	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Defense Documentation Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Naval Ocean Systems Center Attn: Dr. S. Yamamoto, Marine Sciences Division San Diego, California 91232	1
Dr. Fred Saalfeld Chemistry Division Naval Research Laboratory Washington, D.C. 20375	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1